

Effectiveness research on psychosocial interventions among high-utilising patients in primary care

Robert L Woolfolk, Lesley A Allen

Department Psychology, Princeton University, Princeton, New Jersey, USA

WHAT IS ALREADY KNOWN ON THIS TOPIC?

Many patients who experience significant health anxiety and somatisation, and who also over-utilise medical services, are unresponsive to standard medical care.¹ Cognitive behavioural therapy (CBT) has been shown to provide modest reductions in symptomatology and health-care utilisation in these patients.¹ Relaxation training (RT) also has been shown to benefit this population, but has been less widely studied.¹

WHAT DOES THIS PAPER ADD?

- ▶ Given that somatising patients often fail to pursue mental health referrals, stepped-care approaches to treatment within primary care have been suggested. Stepped-care approaches provide a very brief intervention to all patients and a more intensive treatment to the subset of patients who are unresponsive to the initial brief intervention. Barsky and colleagues' study extended the scope of previous research on treatment of somatising patients¹ by examining the effectiveness of a stepped-care cognitive-behavioral approach administered by primary care clinicians in a primary care treatment setting.
- ▶ This is the first direct comparison, conducted in primary care, of the clinical effectiveness of cognitive behavioural therapy and relaxation training for easing hypochondriacal beliefs and somatisation symptoms.

LIMITATIONS

- ▶ Although the study examined a group of patients with high rates of healthcare utilisation, the patients were relatively high-functioning compared with those examined in earlier research.² It is not clear that all the patients treated were either high in somatisation or hypochondriasis. Fewer than one-third of randomised participants met Diagnostic and Statistical Manual-IV criteria for hypochondriasis and their mean Whitely Index score and Somatic Symptom Inventory score were lower than those reported in previous trials.³ High-utilising patients who were neither somatising nor hypochondriacal would not be expected to benefit from the interventions examined.

- ▶ The study design did not include a no-treatment or treatment-as-usual control group.
- ▶ Many of the conclusions of the study are based on correlational data and a posteriori inferences, rather than confirmation of a priori hypotheses.

WHAT NEXT IN RESEARCH?

The outcome of primary interest for high-utilising patients would be change in healthcare utilisation. Much larger datasets are required to make meaningful comparisons of healthcare costs. Also, whether primary care clinicians can be trained to administer psychosocial treatments as effectively as specialists with more intensive training is yet to be examined.

COULD THESE RESULTS CHANGE YOUR PRACTICES AND WHY?

This study bolsters the conclusions of other research indicating that brief psychosocial treatments administered in primary care can benefit patients who are high utilisers of medical services.⁴ The study serves as an important reminder of the potential benefits of relaxation training for somatising patients. Given that we employ similar methods to those studied in this research, the findings of this study would not alter our approach to treatment.

Competing interests None.

doi:10.1136/eb-2013-101678

REFERENCES

1. **Allen LA**, Escobar JI, Lehrer PM, *et al.* Psychosocial treatments for multiple unexplained physical symptoms: a review of the literature. *Psychosom Med* 2002;**64**:939–50.
2. **Allen LA**, Woolfolk RL, Escobar JI, *et al.* Cognitive-behavioral therapy for somatization disorder: a randomized controlled trial. *Arch Intern Med* 2006;**166**:1512–18.
3. **Barsky AJ**, Ahern DK. Cognitive behavior therapy for hypochondriasis: a randomized controlled trial. *JAMA* 2004;**291**:1464–70.
4. **Escobar JI**, Gara MA, Diaz-Martinez AM, *et al.* Effectiveness of a time-limited cognitive behavior therapy type intervention among primary care patients with medically unexplained symptoms. *Ann Fam Med* 2007;**5**:328–35.

ABSTRACT FROM: Barsky AJ, Ahern DK, Bauer MR, *et al.* A randomized trial of treatments for high-utilizing somatizing patients. *J Gen Intern Med* 2013;**70**:1396–404.

Participants Eighty-nine people selected from the highest 20% of healthcare utilisers in the prior year, identified through encounter claims data, and whose names were approved by their clinicians. Potential participants completed a telephone screening questionnaire to measure levels of medically unexplained symptoms and health-related anxiety. Those who scored above a pre-determined cut-off were eligible to participate. Participants were excluded if they had more than one visit to the same subspecialty or more than one narcotic prescription during the previous year, a diagnosis of alcoholism or psychosis, symptom-contingent litigation or disability compensation or were under

18 or over 70 years of age. Less than one-third of participants met Diagnostic and Statistical Manual-IV (DSM-IV) criteria for hypochondriasis.

Setting Six primary care suites in two large, multispecialty community practices in the USA.

Intervention Cognitive behavioural therapy (CBT, n=59) versus relaxation therapy (RT, n=30). CBT consisted of two steps. The first step was of three, manualised 60 min counselling sessions focusing on symptom amplification, cognitive distortions and misunderstandings about symptoms, disease and medical care. Participants judged to have not improved after this step had a