
Evidence-Based Mental Health



Contents

EBMH Notebook

The STAR*D trial: the 300 lb gorilla is in the room, but does it block all the light? Simon Hatcher

Unravelling factor analysis Khalida Ismail 99

102 Et al

Prognosis

- Partial syndromes of eating disorders in adolescence are associated with adverse health outcome in adulthood
- Risk of death from natural causes increased in people with psychiatric admissions

Diagnosis

107 Common mental disorders in primary care in India: little to distinguish five commonly used screening instruments

Therapeutics



Review: Individual behavioural therapy reduces symptoms of depression

- 109 Mood and anxiety disorders: differences between long- and short-term psychotherapies in work ability, but not employment
- **110** Adolescents with SSRI-resistant depression: CBT plus antidepressant switch more effective than medication switch alone
- Adjunctive aripiprazole improves symptoms in antidepressant refractory major depressive disorder
- 112 Review: Insufficient evidence on Huperzine A for Alzheimer's
- 113 Review: Cholinesterase inhibitors and memantine consistently but marginally improve symptoms of dementia
- 114 Review: Paliperidone reduces symptoms of schizophrenia and schizophrenia-like illness
- Second generation antipsychotics reduce treatment discontinuation rates compared with haloperidol

November 2008 Vol 11 No 4

- Little difference between haloperidol, risperidone and olanzapine in weight gain after 1 year of treatment
- **117** Review: Cognitive remediation improves cognitive functioning in schizophrenia
- 118 Transdermal methylphenidate more effective than placebo for treating ADHD
- A parenting programme does not affect behavioural problems in toddlers
- 120 Borderline personality disorder: STEPPS improves symptoms
- Carers of people with dementia: no benefit of adding befriending initiative to usual care
- Timely encouraging letters increase attendance at first consultations at outpatient psychiatric clinics
- Review: Substance use disorders respond to psychosocial interventions
- Review: Short stay hospitalisation does not increase readmissions compared with long stay

Aetiology

- Use of antidepressants in pregnancy does not increase risk of postpartum haemorrhage
- Deployment with combat exposure increases the risk of new-onset PTSD
- Maternal exposure to death of a first degree relative during first trimester of pregnancy increases risk of schizophrenia in offspring

Prevalence



Global WHO survey: poor physical and mental health more prevalent among women who have experienced intimate partner violence

Electronic pages

Purpose and procedure



This article has been chosen by the Editor to be of special interest or importance and is freely available online.



 $\mathbf{C} \left| \mathbf{O} \right| \mathbf{P} \left| \mathbf{E} \right|$ committee on publication ethics

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk

