


Evidence-Based Mental Health



Contents

August 2011 Vol 14 No 3

Purpose and procedure

- 57 Purpose and procedure

EBMH Notebook

- 59 Risk assessment: predicting violence

Editorial

- 62 Ethics of evidence

Aetiology

- 63 Review: childhood ADHD increases the risk of nicotine use in adolescence and alcohol use in young adulthood
- 64 Girls who have bullied others by age 8 are more likely to have a child when they are teenagers
- 65 Increased risk of psychotic symptoms at 12 years in children who have been maltreated by adults or bullied by peers
- 66 Biological parent suicide and severe psychiatric morbidity are risk factors for suicide in adopted and non-adopted offspring
- 67 Use of antidepressants for 12 weeks or longer is associated with a reduced incidence of heart attack in depressed adults

Prevalence

- 68 Regional variation in antipsychotic polypharmacy for schizophrenia

Prognosis

- 69 People with psychotic disorders are at greater risk of conviction for arson
- 70 Incident cannabis use in adolescents and young adults is associated with an increased risk of developing psychotic symptoms

Review

- 71 Sequential use of pharmacotherapy and psychotherapy reduces rates of relapse or recurrence in patients with major depressive disorder

Therapeutics

- 72 Short-term treatments may not affect longer term recovery or recurrence among adolescents with a major depressive episode

- 73 Inhaled loxapine reduces acute agitation in people with schizophrenia compared with placebo
- 74 'Timid to Tiger' group parenting training reduces anxiety diagnoses in 3–9-year-olds
- 75 Bright light treatment reduces symptoms in older adults with non-seasonal major depression
- 76 Review: limited evidence for use of second-generation antipsychotics in anxiety disorders
- 77 Problem-solving therapy reduces disability more than supportive therapy in older adults with major depression and executive dysfunction
- 78 'Tuning in to Kids' parenting programme improves parental emotional awareness, and child behaviour and emotional knowledge
- 79 Effect of first-generation perphenazine and second-generation antipsychotics on depressive symptoms in schizophrenia: all antipsychotics improved symptoms; quetiapine was superior to risperidone for people with major depression at baseline
- 80 Review: antidepressants are safe and effective for people with depression who have additional chronic physical health problems
- 81 Secondary prevention of cardiovascular disease: CBT, in combination with standard care, is effective
- 82 Review: benefits of provision of general physical health advice for people with serious mental illness unclear
- 83 Adding 8 weeks of behavioural therapy to atomoxetine does not improve classroom behaviour in children with ADHD
- 84 Adding group psychotherapy to routine care does not improve outcomes in adolescents who repeatedly self-harm
- 85 Most adolescents achieve longer term remission from major depressive disorder regardless of type of short-term second-line treatment


Treatment

- 86 Review: compulsory community treatment does not improve social functioning or service use in people with severe mental illness

Corrections

- 88 Corrections

Glossary

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org

 equator network

 recycle

When you have finished with this please recycle it