
Evidence-Based Mental Health



Contents

Editorial

Time to abandon Evidence Based Medicine?

Expert review

Bipolar disorder and disruptive mood dysregulation in children and adolescents: assessment, diagnosis and treatment

Aetiology

- Depression and use of antidepressants is associated with increased risk of Clostridium difficile infection
- When viewing empathy-eliciting scenarios, incarcerated men with high psychopathy display differences in brain activity compared with those with low psychopathy
- Adults with psychosis or psychotic experiences may report adverse life events prior to the onset of symptoms but direction of causality cannot be inferred
- The association between maternal depression and anxiety during pregnancy and child attention problems may be partly explained by postnatal symptoms

Children & Adolescents

- Vulnerability for psychosis at ages 13 and 16 predicts cannabis use at ages 16 and 19, and cannabis use at age 16 predicts psychosis vulnerability at age 19
- Personality-targeted interventions delivered by 100 teachers may be effective at reducing alcohol use
- Experiencing maltreatment during childhood increases the likelihood of lifetime alcohol dependence
- Exposure to intimate partner violence and parental depression increases risk of ADHD in preschool children
- Non-transient childhood conversion disorder is uncommon in the UK and Ireland, but use of services is high
- Methylphenidate reduces ADHD symptoms in children with severe ADHD and intellectual disability
- Limited evidence that a brief education programme for parents of high-risk preschool children may reduce risk of internalising disorders in adolescence in girls but not boys
- Suicidal behaviours are common among US 106 adolescents and are associated with mental health disorders

November 2013 Vol 16 No 4

Medical comorbidity

- Centralised depression care according to patient preference is effective for the treatment of depression symptoms following acute coronary syndrome
- Review: insufficient evidence to recommend routine screening for psychological distress among people with cancer
- Current use of second-generation antipsychotics may increase risk of pneumonia in people with schizophrenia
- Review: depression after myocardial infarction is associated with increased risk of all-cause mortality and cardiovascular events

Pharmacotherapy

- Limited evidence that benzodiazepines are beneficial in the treatment of disturbed behaviour
- Review: lithium reduces the risk of suicide 112 compared with placebo in people with depression and bipolar disorder

Prevalence

Review: about one in five people with first-episode psychosis have a history of deliberate self-harm



Family history of REM sleep behaviour disorder more common in individuals affected by the disorder than among unaffected individuals

Psychosocial

- **115** A false-positive on screening mammography has a negative psychosocial impact up to 3 years after receiving the all clear
- Community treatment orders do not reduce hospital readmission in people with psychosis
- Letter in response to Dr Segal's commentary
- Mortality risks from psychological distress are greater for people of lower socioeconomic status

Psychotherapy

- Review: computerised CBT improves adult depression in the short-term, but its effect may have been overestimated previously
- Interpersonal therapy with parenting enhancement does not reduce depression symptoms in low-income mothers compared with control
- Lay therapist-delivered manualised CBT improves anger coping in people with intellectual disability and anger problems, but effects on anger measures mixed
- Review: different psychotherapies have comparable effects on adult depression
- CBT increases remission of child or adolescent anxiety disorders compared with wait list control

