


# Evidence-Based Mental Health



## Contents

### Editorial

- 1 Advancing evidence-based practice to improve patient care

### Psychological interventions

- 3 Review: evidence-based psychotherapies are more effective than usual care for young people with psychopathology: effects are influenced by location of care and participant characteristics
- 4 Antidepressants increase risk of behavioural symptoms in children and adolescents with depression or anxiety
- 5 Brief, personality-targeted, teacher-delivered CBT interventions reduce depression, anxiety and conduct disorder symptoms in high-risk adolescents
- 6 In children treated for internalising or externalising problems, sudden large improvements between sessions is associated with longer term outcomes

### Pharmacological interventions

- 7 Review: psychotherapy, somatic therapy and pharmacotherapy are all more effective than control for the treatment of PTSD
- 8 Review: maintenance antidepressants reduce risk of relapse in the 6 months following ECT in people with major depression
- 9 Review: 15 antipsychotic drugs are more effective than placebo for the treatment of schizophrenia, but vary in their tolerability
- 10 A dose reduction/discontinuation strategy improves long-term recovery in people with remitted first-episode psychosis compared to maintenance therapy

### Statistics in practice

- 11 Exploring and accounting for publication bias in mental health: a brief overview of methods

### Causes

- 16 Review: no evidence that stimulant medication for ADHD influences lifetime risk of substance use or dependence

## February 2014 Vol 17 No 1

- 17 Children conceived following induced ovulation or intrauterine insemination have a small increased risk of mental disorders
- 18 Different genetic factors influence specific symptom dimensions of DSM-IV major depression
- 19 Adolescents with emotional, conduct and hyperkinetic disorders who are experiencing psychotic symptoms may be at increased risk of suicide attempt
- 20 Severe infection and autoimmune disease are associated with increased risk of mood disorders

### Outcomes

- 21 Chronic care management programme is no more effective than usual primary care at increasing abstinence among people with alcohol or substance dependency
- 22 Review: recommendations for the assessment and management of sleep disorders in ADHD
- 23 A social skills and parental training intervention for disruptive boys reduces substance use behaviours in adolescence
- 24 Review: insufficient evidence on the effectiveness of interventions to prevent child sexual abuse in individuals at risk of abusing

### Evidence-based case conference


- 25 Antidepressants during ECT

### Prevalence, assessment and diagnosis

- 27 Over 60% of US adolescents have experienced a potentially traumatic event, almost 8% of whom have associated PTSD
- 28 Review: one in seven homicides worldwide is perpetrated by an intimate partner
- 29 Age may moderate response to different unguided Internet-delivered interventions for depression

### Statistics in pills

- 30 How to assess publication bias: funnel plot, trim-and-fill method and selection models

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

[www.publicationethics.org](http://www.publicationethics.org)

 equator network

 recycle

When you have finished with this please recycle it

Receive regular table of contents by email. Register using this QR code.

