


Evidence-Based Mental Health



Contents

May 2014 Vol 17 No 2

Editorial

- 31 Nulla dies sine linea

Clinical review

- 33 Pharmacotherapy of treatment-resistant schizophrenia: a clinical perspective



Causes

- 38 Prenatal and postnatal exposure to an unhealthy diet is associated with behavioural and emotional problems in children
- 39 In people with ultra high risk symptoms, risk of transition to psychotic disorders is highest in the first 2 years

Outcomes

- 40 First randomised controlled-trial research on seclusion and restraint reduction achieves intent
- 42 The outcomes of a school-based intervention for depressive symptoms in adolescents do not echo the promising findings of earlier studies
- 43 Recovery from first-episode psychosis: early identification and prompt initiation of treatment improve long-term outcomes

Evidence-based case conference

- 45 Diagnostic test for dementias

Prevalence, assessment and diagnosis

- 47 Effectiveness research on psychosocial interventions among high-utilising patients in primary care
- 49 In young men, various risk factors are associated with later development of young-onset dementia

Pharmacological interventions

- 50 Longer term outcomes need to be examined for enhancing the treatment of social anxiety disorder after insufficient improvement with sertraline

- 51 Metformin for weight loss in schizophrenia: safe but not a panacea

Statistics in practice

- 53 Demystifying fixed and random effects meta-analysis

Psychological interventions

- 58 Augmentation with cognitive behavioural therapy has superior efficacy to augmentation with risperidone for treating adults with treatment-resistant OCD
- 59 Continuation-phase cognitive therapy and fluoxetine are effective in reducing the risk of relapse/recurrence in major depression after incomplete remission

Miscellaneous

- 61 To disclose or not: does a decision tool help?
- 62 Financial incentives improve adherence to maintenance antipsychotic treatment

Statistics in pills


- 64 How to interpret meta-analysis models: fixed effect and random effects meta-analyses

PostScript

- 48 Corrections

Online articles

- e1 Sex disparities in depression prevalence are lost when male-type depression symptoms are considered alongside traditional symptoms
- e2 Behavioural treatment combined with buprenorphine does not reduce opioid use compared with buprenorphine alone
- e3 Review: psychosocial mood management improves smoking cessation in people with current and past depression

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org

 equator network

 recycle

When you have finished with this please recycle it

Receive regular table of contents by email. Register using this QR code.

