


Evidence-Based Mental Health



Contents

November 2014 Vol 17 No 4

Clinical review

- 97 Self-harm in young people



Causes and risk factors

- 100 Young maternal age and old paternal age induce similar risk of mental disorders in offspring
- 101 High rates of suicide attempt in early-onset psychosis are associated with depression, anxiety and previous self-harm
- 102 Medication reduces crash risk among men with ADHD

Evidence-based case conference

- 103 Adverse effects of antidepressants during pregnancy



Prevalence, assessment and diagnosis

- 105 Developing accuracy when estimating Global Burden of Disease (GBD) in mental and substance use disorders using complex research methodology, refined data analysis and community participation
- 106 Neonatal factors associated with autism spectrum disorders in infants
- 107 Children exposed to trauma should be screened for symptoms of PTSD

Pharmacological interventions

- 108 Atomoxetine is a second-line medication treatment option for ADHD
- 109 Ziprasidone monotherapy is ineffective for bipolar-associated anxiety
- 110 Paliperidone depot is not different from haloperidol in relapse prevention of schizophrenia, but different side effects should be considered

Statistics in practice

- 111 A hands-on practical tutorial on performing meta-analysis with Stata




Psychological interventions

- 117 Extending the evidence base for prolonged exposure: significant effectiveness noted with sexual abuse-related PTSD in adolescent girls
- 118 Finally moving beyond the horse race: CBT and psychodynamic therapy equally effective for depression
- 119 Some evidence of efficacy but CBT not the answer to mental health needs in general hospitals

Miscellaneous

- 120 Theory of mind interventions can be effective in treating autism, although long-term success remains unproven
- 121 Targeted, preventive interventions in primary school prevent mental health problems in adolescence
- 122 Sleep therapy is effective in reducing symptoms of depression among adolescents
- 123 Review tempers expectations for peer-led services but further studies are required
- 124 Community-based intervention for people with schizophrenia successfully provided by supervised community health workers in a low-resource setting

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org

 equator
network

 recycle

When you have finished with this please recycle it

Receive regular table of contents by email. Register using this QR code.

