

Evidence-Based Mental Health

Contents

Purpose and procedure (abbreviated)	34	Review: quetiapine leads to fewer withdrawals because of adverse effects in schizophrenia	52
EBMH notebook		Paroxetine reduced symptoms and avoidance in generalised social phobia	53
Asking structured and focused clinical questions: essential first step of evidence-based practice	35	Diagnosis	
Mental health technology assessment: practice based research to support evidence-based practice.....	37	An informant questionnaire and a standard cognitive test improved screening for dementia	54
Other articles noted	40	The Alcohol Use Disorders Identification Test was not affected by ethnic and sex bias	55
Therapeutics		Prognosis	
Review: self help interventions alone minimally increase smoking cessation	41	Apolipoprotein E ϵ 4 allele predicted an accelerated rate of cognitive decline in Alzheimer's disease	56
Review: donepezil improves clinical state in mild or moderate Alzheimer's disease	42	One year after head injury a substantial proportion of patients showed neuropsychiatric symptoms	57
Cognitive behavioural therapy was effective for adolescent depression after controlling for adverse predictors of outcome	43	Personality traits and coping styles were associated with depression in spinal cord injury.....	58
CBT for major depression may continue to reduce relapse in the long term	44	Children and adolescents with severe traumatic brain injury had verbal encoding deficits at 1 and 3 months after the injury	59
CBT was effective at 2 years in recurrent depression ...	45	Aetiology	
Review: a lack of evidence exists to indicate the most effective treatment for patients who have deliberately harmed themselves	46	Review: incidence of dementia and Alzheimer's disease increases with increasing age	60
Review: thioridazine is not more effective than placebo or other neuroleptic drugs in elderly patients with dementia	47	Use of benzodiazepines increased road traffic accidents whereas use of tricyclic antidepressants and SSRIs did not	61
Haloperidol, 2–3 mg/day, decreased psychosis and disruptive behaviours in Alzheimer's disease	48	Adverse perinatal conditions were associated with risk of suicide by violent means for men	62
Review: St John's wort is more effective than placebo for treating depressive disorders	49	Genetic risk factors influenced cocaine abuse and dependence more than cocaine use in women.....	63
Cognitive behaviour therapy relieved depression in type 2 diabetes mellitus	50	Cognitive function during early childhood did not improve with declining blood lead concentrations.....	64
Cognitive behavioural therapy reduced cocaine abuse compared with 12 step facilitation	51	Glossary	Inside back cover