Evidence-Based Mental Health



Contents

Editorial

Providing the most appropriate care to our individual patients

A Cipriani, A Tomlinson

Original articles

Understanding the quality, effectiveness and attributes of top-rated smartphone health apps

H Wisniewski, G Liu, P Henson, A Vaidyam, N K Hajratalli, J-P Onnela, J Torous

Development and validation of a prediction model for the probability of responding to placebo in antidepressant trials: a pooled analysis of individual patient data

> K Shinohara, S Tanaka, H Imai, H Noma, K Maruo, A Cipriani, S Yamawaki, T A Furukawa

Clinical reviews

Digital technology for health promotion: opportunities to address excess mortality in persons living with severe mental disorders J A Naslund, K A Aschbrenner

What is the evidence for the use of simulation training to teach communication skills in psychiatry?

J Neale

Systematic reviews

26 Comparative evaluation of group-based mindfulness-based stress reduction and cognitive behavioural therapy for the treatment and management of chronic pain: A systematic review and network meta-analysis

> E-L Khoo, R Small, W Cheng, T Hatchard, B Glynn, D B Rice, B Skidmore, S Kenny, B Hutton, P A Poulin

February 2019 Volume 22 Issue 1

36 Effects of mindfulness-based interventions on alexithymia: a systematic review

H Norman, L Marzano, M Coulson, A Oskis

Statistics in practice

Reasoning under uncertainty



C Aitken, D Mavridis

Electronic pages

Attention-deficit/hyperactive disorder: missing the bull's eye

C Leon, R Sharma, S Kaur

Reply to Leon et al S Cortese, D Coghill

Adjunctive minocycline in schizophrenia: what one well-conducted study can tell us (and what it can't)

M Solmi, C U Correll

CBT, medication and the combination are effective for childhood anxiety

L M Hana, E McIngvale, M Davis, E A Storch

e5 Electroconvulsive therapy does not increase the risk of dementia in patients with affective disorders

P Sienaert

There are promising effects from group intervention for clinically aggressive youth on probation

A Dopp

Preventive cognitive therapy could be a viable and effective addition to antidepressant medication in preventing relapse or recurrence in major depressive disorder

M J Huijbers, C Wentink, A E M Speckens



This article has been chosen by the Editor to he of special interest or importance and is freely available online.



This article has been made freely available OPEN ACCESS online under the BMJ Journals open access scheme. See http://ebmh.bmj.com/site/ about/guidelines.xhtml#open



This journal is a member of and subscribes to the principles of the Committee on Publication Ethics www.publicationethics.org





ve regular table of contents by email.



ebmh.bmj.com