

Evidence-Based Mental Health

Contents

Purpose and procedure	2	Once a day Concerta methylphenidate was equivalent to 3 times daily methylphenidate in children with ADHD	20
EBMH notebook		Quintuple the daily maintenance dose of buprenorphine given every 5 days was associated with opioid withdrawal symptoms	21
The scientific evaluation of mental health treatments: an historical perspective	4	Repeated doses of porcine secretin did not improve symptoms, language, or cognitive functioning in children with autism or autism spectrum disorder	22
Evidence-based practice and clinical social work	6	Cognitive treatment reduced symptoms and improved functioning in bipolar disorder	23
Other articles noted	8	St John's wort was not better than placebo for reducing depression scores	24
Therapeutics		Paroxetine and sertraline delayed ejaculation in heterosexual men with lifelong rapid ejaculation	25
A geriatric evaluation and management programme prevented functional decline and reduced depression in high risk older adults	9	Diagnosis	
Lithium is more effective than placebo for preventing all relapses in patients with bipolar but not unipolar disorder	10	Review: most laboratory tests do not add to the diagnostic accuracy of clinical criteria for dementia	26
Risperidone was effective for aggression in adolescents with disruptive behaviour disorders and below average intelligence	11	Clinical Prediction Guide	
Life skills training and booster sessions plus drinking reduction treatment was effective in heavily drinking women	12	A risk assessment system based on preoperative factors was fairly specific in predicting postoperative delirium risk in patients having major elective surgery	27
Review: eye movement desensitisation and reprocessing is not better than exposure therapies for anxiety or trauma	13	Qualitative	
Brief psychodynamic interpersonal therapy after deliberate self poisoning reduced suicidal ideation and deliberate self harm	14	A 4 stage model described how clients and their relatives experienced the process of developing schizophrenia ..	28
Review: placebo is better than no treatment for subjective continuous outcomes and for treatment of pain	15	Parents' experiences of having children with a significant developmental disability	29
Paroxetine but not imipramine improved depression related outcomes in adolescents with <i>DSM-IV</i> major depression	16	Drug users' experiences of injecting included descriptions of needle fixation	30
Review: iron treatment does not improve psychomotor development and cognitive function at 30 days in children with iron deficiency anaemia	17	General practitioners often used emotional responses for "referrals away" and intellectual decision making for "referrals to" in patients with mild depression and/or anxiety	31
Olanzapine was more effective than lorazepam at 2 hours but not at 24 hours in bipolar mania with acute agitation	18	Caregivers described how an Alzheimer's disease respite programme gave them time to attend to their own needs	32
Ongoing contact for 2 years after discharge was associated with greater survival in patients at risk of post crisis suicide	19	Glossary	Inside back cover