

# Evidence-Based Mental Health

## Contents

### EBMH Notebook

What are the effects of drug treatments for panic disorder? . . . . .	34
Difficulties developing evidence-based approaches in learning disabilities . . . . .	37

### Resource Corner . . . . .

### Therapeutics

Vitamin and fatty acid supplements may reduce antisocial behaviour in incarcerated young adults . . . . .	41
Once-daily atomoxetine may reduce attention deficit hyperactivity disorder symptoms in children and adolescents . . . . .	42
Review: adderall may have a small advantage over methylphenidate in attention deficit hyperactivity disorder . . . . .	43
20mg transdermal selegiline daily may be effective and well tolerated in adults with major depression . . . . .	44
Review: antidepressants and psychotherapy may be equally effective for promoting remission in major depressive disorder . . . . .	45
Review: Low dose tricyclic antidepressants may be effective for adults with acute depressive disorder . . . . .	46
Enhanced primary care may encourage remission in depression . . . . .	47
Review: there is limited evidence about the effectiveness of interventions for treatment-refractory depression . . . . .	48
18-month maintenance treatment with sertraline may have sustained psychosocial benefits in chronic depression . . . . .	49
Skills training plus exposure therapy may reduce post traumatic stress in women who experienced childhood abuse . . . . .	50
Fluoxetine may prevent relapse in post traumatic stress disorder . . . . .	51
Stress management programme may reduce hospital admissions among people with schizophrenia . . . . .	52

Review: there is no strong evidence to support antipsychotic combination therapy in schizophrenia . . . . .	53
Cognitive training may improve targeted cognitive functions in older adults . . . . .	54
Review: cognitive behavioural interventions may be effective for chronic fatigue syndrome and chronic back pain . . . . .	55
Group interpersonal psychotherapy may be as effective as group cognitive behavioural therapy for overweight people with binge eating disorder . . . . .	56
Review: there is insufficient evidence for naltrexone maintenance treatment in opioid dependence . . . . .	57
Skills training programmes for significant others may encourage unmotivated drug users to seek treatment . . . . .	58

### Prevalence

Almost half of people suffering traumatic brain injury may later be diagnosed with axis I disorders . . . . .	
---	--

### Aetiology

Depressive symptoms in adolescence may increase the risk of psychiatric disorders in early adulthood . . . . .	60
Individual and familial mental illness is associated with an increased risk of suicide in young people . . . . .	61
There is little evidence that combined vaccination against measles, mumps, and rubella is associated with autism . . . . .	62

### Diagnosis

Case-finding instruments may help identify depression in primary care . . . . .	63
---	----

### Qualitative

Many factors make family members caring for an elderly relative with dementia reluctant to use support services . . . . .	64
---	----