

Evidence-Based Mental Health

Contents

EBMH Notebook

The internet and mental health practice	66
Which treatments are effective for cognitive, behavioural and psychological symptoms in dementia?	69
The number needed to treat: problems describing non-significant results	72

Prevalence

About 3 in every 1000 US children in large metropolitan areas may have autism or related developmental disorders ...	73
Review: lifetime prevalence of schizophrenia and related disorders is about 5.5 per 1000, but there is significant variation between regions.	74
Review: there is marked socioeconomic inequality in persistent depression	75
Review: black people are more likely than white people to be detained in psychiatric wards in the United Kingdom ...	76

Diagnosis

Review: antenatal screening tools to predict postnatal depression generally have low specificity and sensitivity	77
Local adaptation of guidelines for identifying mental disorders in primary care may not improve patient outcomes.	78

Therapeutics

Review: atypical antipsychotics and psychosocial interventions, alone or in combination, may reduce youth aggression.	79
Review: it is unclear whether treating internalising disorders in childhood prevents substance abuse in adolescence	80
Review: parent training and community-based interventions may benefit children with disruptive behaviour disorders	81
Problem solving programme implemented by teachers may prevent depression in the short term, but longer term benefits are unclear.	82

Review: electroconvulsive therapy may be an effective short term treatment for people with depression.	83
---	----

Review: continuing antidepressants for 12 months following acute treatment may reduce the risk of relapse in depression	84
---	----

Adjunctive cognitive therapy is more effective, but more costly, than standard clinical management for relapse prevention in depression	85
---	----

Collaborative care may improve depression management in older adults	86
--	----

Review: counselling in primary care may have short term benefits for people with common mental disorders.	87
--	----

Advance directives about treatment preferences had little impact on compulsory readmissions for people with serious mental illness	88
--	----

Review: women with schizophrenia have poorer pregnancy outcomes than other women, but it is unclear whether antipsychotic medications affect their infants.	89
--	----

The benefits of exposure therapy alone may last longer than sertraline alone or sertraline plus exposure therapy in social phobia	90
---	----

Hydroxyzine may be safe and effective in generalised anxiety disorder	91
---	----

Buprenorphine may be more effective than clonidine and other symptomatic medications for short term ambulatory heroin withdrawal	92
--	----

Clozapine may be more effective than olanzapine for reducing suicidal behaviour in people with schizophrenia at high risk	93
---	----

Review: cholinesterase inhibitors have a modest effect on neuropsychiatric and functional outcomes in Alzheimer's disease	94
---	----

Aetiology

Ethnicity may be linked to thin body preoccupation and social pressure in the development of eating disorders	95
--	----

Qualitative

People with schizophrenia believe that they are stigmatised at work and in the community	96
--	----