Evidence-Based Mental Health

Contents

EBMH Notebook

The internet and mental health practice 66

Which treatments are effective for cognitive, behavioural and psychological symptoms in dementia? 69

The number needed to treat: problems describing non-significant results 72

Prevalence

About 3 in every 1000 US children in large metropolitan areas may have autism or related developmental disorders ... 73

Review: lifetime prevalence of schizophrenia and related disorders is about 5.5 per 1000, but there is significant variation between regions 74

Review: there is marked socioeconomic inequality in persistent depression 75

Review: black people are more likely than white people to be detained in psychiatric wards in the United Kingdom ... 76

Diagnosis

Review: antenatal screening tools to predict postnatal depression generally have low specificity and sensitivity 77

Local adaptation of guidelines for identifying mental disorders in primary care may not improve patient outcomes 78

Therapeutics

Review: atypical antipsychotics and psychosocial interventions, alone or in combination, may reduce youth aggression 79

Review: it is unclear whether treating internalising disorders in childhood prevents substance abuse in adolescence 80

Review: parent training and community-based interventions may benefit children with disruptive behaviour disorders 81

Problem solving programme implemented by teachers may prevent depression in the short term, but longer term benefits are unclear 82

Review: electroconvulsive therapy may be an effective short term treatment for people with depression 83

Review: continuing antidepressants for 12 months following acute treatment may reduce the risk of relapse in depression 84

Adjunctive cognitive therapy is more effective, but more costly, than standard clinical management for relapse prevention in depression 85

Collaborative care may improve depression management in older adults 86

Review: counselling in primary care may have short term benefits for people with common mental disorders 87

Advance directives about treatment preferences had little impact on compulsory readmissions for people with serious mental illness 88

Review: women with schizophrenia have poorer pregnancy outcomes than other women, but it is unclear whether antipsychotic medications affect their infants 89

The benefits of exposure therapy alone may last longer than sertraline alone or sertraline plus exposure therapy in social phobia 90

Hydroxyzine may be safe and effective in generalised anxiety disorder 91

Buprenorphine may be more effective than clonidine and other symptomatic medications for short term ambulatory heroin withdrawal 92

Clozapine may be more effective than olanzapine for reducing suicidal behaviour in people with schizophrenia at high risk 93

Review: cholinesterase inhibitors have a modest effect on neuropsychiatric and functional outcomes in Alzheimer's disease 94

Aetiology

Ethnicity may be linked to thin body preoccupation and social pressure in the development of eating disorders 95

Qualitative

People with schizophrenia believe that they are stigmatised at work and in the community 96

BMJ Ment Health: first published as on 1 August 2003. Downloaded from http://mentalhealth.bmj.com/ on April 27, 2024 by guest. Protected by copyright.