

Evidence-Based Mental Health

Contents

Purpose and procedure	2	A structured needs assessment does not improve clinical outcomes for clients under the care programme approach.....	18
Prognosis			
Chronic, long episode mania exists in children	5	An individual placement and support programme is more effective than skills training and temporary employment for people with severe mental illness.....	19
Review: a pooled long term persistence rate of 40% for childhood OCD is lower than previously expected	6		
Therapeutics		Aetiology	
Intramuscular haloperidol-promethazine sedates violent or agitated patients more quickly than intramuscular lorazepam.....	7	Eating disorders, especially anorexia nervosa, are associated with an increased risk of attempted suicide in young women	20
Cognitive behavioural therapy reduces short term rehospitalisation compared with psychoeducation in inpatients with schizophrenia.....	8	Review: depression and anxiety during pregnancy are strong indicators of postpartum depression	21
Adding psychosocial therapy to methylphenidate may not improve its effectiveness in stimulant responsive children with ADHD	9	Review: tricyclic related drugs, but not classical tricyclics, are a tolerable alternative to SSRIs in older people with depression.....	22
Fluoxetine plus cognitive behavioural therapy improves symptoms of major depressive disorder in adolescents	10	Review: vaccines containing thimerosal are not associated with autistic spectrum disorders in children.....	23
School-based interpersonal psychotherapy improves depression in older adolescents.....	11	Stable monotherapy with clozapine or olanzapine increases the incidence of diabetes mellitus in people with schizophrenia	24
Review: long term psychotherapy in combination with antidepressants increases compliance and response rates in people with depression.....	12	Prevalence	
St John's Wort improves somatoform disorders	13	Major depression is common in people over the age of 50, particularly in those at socioeconomic disadvantage, or with poor physical health and previous depressive symptoms.....	25
Review: acamprosate and naltrexone are safe and effective but have low compliance rates for people with alcohol dependence	14	Canadian study finds that antidepressant use has increased in people with major depression over the past decade	26
Review: donepezil improves cognitive and global function in people with mild to moderate Alzheimer's disease.....	15	PTSD plus depression are comorbid conditions but depression can occur independently in the acute aftermath of trauma	27
Review: atypical antipsychotics may be useful in treating behavioural and psychological symptoms of dementia but cause adverse effects.....	16	Mental disorders are highly prevalent in Germany	28
A joint crisis plan negotiated with mental health staff significantly reduces compulsory admission and treatment in people with severe mental illness	17	Miscellaneous	
		Other articles noted	3