Evidence-Based Mental Health

Contents

Purpose and procedure	58	Cognitive behavioural therapy has short term but not long term benefits in people with residual symptoms of depression	75
EBMH Notebook		uepression	15
Rethinking evidence-based practice for children's mental health	60	Improving quality of primary care reduces depression and improves quality of life in adolescents	76
bmjupdates ⁺ , a new FREE service for evidence-based clinical practice	62	Review: cognitive behaviour therapy is beneficial in children with anxiety disorders	77
Other articles noted	64	Adding mixed amphetamine salts to divalproex sodium improves ADHD symptoms in children with bipolar disorder and comorbid ADHD	78
Prognosis		Valproate reduces alcohol consumption in people with	
Functional impairment recovers after episodes of major depression	65	comorbid alcohol dependency and bipolar disorder	79
Review: depression may increase mortality in coronary		Lithium may reduce gambling severity in pathological gamblers with bipolar disorder	80
heart disease	66	Adding a nurse-based intervention programme to usual	
Review: after myocardial infarction, depression and poor prognosis are associated	67	care improves manic symptoms in people with bipolar disorder	81
People diagnosed with schizophrenia retain the diagnosis long term	68	Glycine and D-serine improve the negative symptoms of schizophrenia	82
Diagnosis		Review: pharmacotherapy provides minimal improvements in the neuropsychiatric symptoms of	
The Screening Tool for Autism in Two Year Olds can identify children at risk of autism	69	dementia	83
Middelheim Frontality Score may be useful for	09	Review: short term psychotherapy is an effective treatment for psychiatric disorders	84
differentiating between Alzheimer's disease and frontotemporal dementia	70	Astialame	84
The control of the		Aetiology	
Therapeutics		Review: migrants are at increased risk of developing schizophrenia	85
Review: hypericum extracts are safer and lead to fewer adverse effects than older standard antidepressants, but have similar tolerability to SSRIs	71	Low intelligence increases risk of suicide	86
Metyrapone is an effective adjuvant treatment for major depression	72	Cannabis use increases the risk of young people developing psychotic symptoms, particularly if already predisposed	
Bright light treatment an effective addition to antidepressants for non-seasonal depression	73	Prevalence	87
Prophylactic mirtazapine may help to prevent post- stroke depression in people with good cognitive	74	Intoxicant use increased among Norwegian adolescents between 1992 and 2002, and attempted suicides increased among girls	88