

# Evidence-Based Mental Health

## Contents

<b>Purpose and procedure (abbreviated)</b> .....	98
<b>EBMH notebook</b>	
Evaluating qualitative research .....	99
Evidence-based medicine and complementary medicine .....	102
<b>Other articles noted</b> .....	104
<b>Therapeutics</b>	
A very brief 3 session psychotherapy intervention was effective in subsyndromal depression .....	105
Review: lower doses of antidepressant drugs are effective and have fewer adverse effects in depression ..	106
Intravenous lorazepam reduced alcohol related seizures in patients with chronic alcohol abuse .....	107
Equivalent response to methylphenidate in children with ADHD plus anxiety and those with ADHD only ..	108
Slow release morphine did not reduce neonatal abstinence syndrome more than methadone .....	109
Brief physician advice reduced drinking in older adults ..	110
Coping skills training was effective in adolescents with type 1 diabetes .....	111
Eptastigmine led to cognitive, clinical, and functional benefits in Alzheimer's disease .....	112
Memantine led to functional improvement and reduced care dependence in severe dementia .....	113
Slow, repetitive transcranial magnetic stimulation was effective for major depression .....	114
Review: greater intensity of light leads to a greater reduction in typical (but not atypical) depressive symptoms in seasonal affective disorder .....	115
Sertraline was effective in children and adolescents with obsessive compulsive disorder .....	116
Cognitive behaviour therapy or temazepam, or both, improved short term outcomes for older adults with persistent insomnia .....	117
Clozapine improved psychosis without worsening parkinsonism .....	118
<b>Diagnosis</b>	
NINCDS-ADRDA criteria were not able to exclude frontotemporal dementia .....	119
<b>Prognosis</b>	
1 year later, 9–13 year olds impaired but not diagnosed were as disturbed as those diagnosed.....	120
Performance on the praxis subscale of the CAMCOG was associated with survival in patients with Alzheimer's disease .....	121
Risk of homelessness was increased in schizophrenia when 3 factors were present .....	122
Most patients returned to previous occupational level 1 year after head injury .....	123
<b>Aetiology</b>	
A decrease in care was associated with suicide in mental illness .....	124
An inverted U relation existed between unemployment and foster home placements .....	125
Current cigarette smoking was associated with a higher risk of Alzheimer's disease .....	126
Older women with depression were at increased risk for falls and fractures .....	127
<b>Economics</b>	
Assertive community treatment for homeless adults with severe mental illnesses was effective but not more expensive than usual care .....	128
<b>Subject index</b> .....	129
<b>Author index</b> .....	134
<b>Contents</b>	
<b>Glossary</b> .....	Inside back cover