

Evidence-Based Mental Health

Contents

Purpose and procedure	2	Oral nalmefene reduced relapse to heavy drinking and was well tolerated in patients with alcohol dependence ..	20
EBMH notebook		A behavioural intervention reduced depression and organic symptoms in elderly people in residential homes	21
Assessing allocation concealment and blinding in randomised controlled trials: why bother?	4	Review: cognitive and behaviour therapies are effective for chronic pain	22
Answering mental health questions with reliable research evidence	6	More people with depression continued treatment with fluoxetine than with desipramine or imipramine	23
Other articles noted	8	Review: nursing interventions increase smoking cessation rates in adults	24
Therapeutics		Prognosis	
A needs based intervention for patients with schizophrenia and their carers reduced relapse and carer needs	9	Failure to thrive in infancy was associated with later stunting and wasting but not cognitive or educational disadvantages	25
Review: rivastigmine may improve cognitive outcomes in Alzheimer's disease	10	Nature and outcome of psychotic illness was similar in all ethnic groups	26
Intensive case management was as effective as standard case management in severe psychotic illness	11	Pattern of course of psychosis and diagnostic stability did not differ by ethnic family background	27
Reframing memory reduced some measures of children's anticipatory distress and pain during lumbar puncture	12	Depressive coping and emotional distress predicted shortened survival in patients with lung cancer	28
Individual drug counselling plus group drug counselling was effective in patients with cocaine dependence	13	Patients with psychiatric disorders, particularly affective disorders, had an increased risk of developing dementia ..	29
Individual placement and support led to more competitive employment than did enhanced vocational rehabilitation for severe mental disorder	14	Aetiology	
Review: studies with sufficient follow up do not show a clear benefit for pharmacotherapy in alcohol dependence	15	Review: schizophrenia is associated with premature rupture of membranes, gestational age < 37 weeks, and use of an incubator or resuscitation	30
Mentally ill adults had fewer homeless days in staffed group homes than in independent flats	16	Typical clinical doses of methylphenidate did not cause tics or exacerbate pre-existing tics in children with attention deficit hyperactivity disorder	31
A literacy promotion intervention increased the frequency of parent to child reading	17	Qualitative	
Cognitive behavioural therapy (CBT) with exercise led to higher levels of continuous smoking cessation in women than did CBT without exercise	18	Women with borderline personality disorder expressed they were living with a pejorative label, with self destructive behaviour viewed as manipulative, and with limited access to care	32
Review: progestogens do not resolve postnatal depression but oestrogens may improve depression scores	19	Glossary	Inside back cover