

# Evidence-Based Mental Health

## Contents

<b>Purpose and procedure</b> .....	34	Involuntary outpatient commitment did not reduce hospital recidivism, though may for some patient subgroups .....	53
<b>EBMH notebook</b>		Integrated outpatient treatment increased abstinence in men with alcohol related illness who were ongoing drinkers .....	54
Evidence based psychotherapy: special case or special pleading? .....	35		
Welcome to our new associate editors .....	37		
A new addition to this journal .....	37		
<b>Resource corner</b> .....	38		
<b>Other articles noted</b> .....	40		
<b>Therapeutics</b>			
Paroxetine was effective for reducing symptoms in social phobia .....	41		
The Fast Track prevention programme improved some outcomes in high risk schoolchildren .....	42		
The Fast Track prevention programme improved classroom atmosphere and peer reported behaviour problems .....	43		
Review: lithium augmentation increases treatment response in refractory depression .....	44		
Sertraline was more effective than desipramine or placebo for premenstrual syndrome .....	45		
Reading and social skills programmes had limited effectiveness on behavioural maladjustment in primary school children .....	46		
A community reinforcement and family training approach was effective in engaging unmotivated problem drinkers in treatment .....	47		
Antidepressants plus cognitive therapy reduced relapse in residual depression .....	48		
Plasma exchange and intravenous immunoglobulin improved symptoms in a subtype of obsessive compulsive disorder and tic disorders .....	49		
Review: psychotherapy may be of benefit in personality disorders .....	50		
Venlafaxine had higher response and remission rates than paroxetine in non-chronic treatment resistant depression .....	51		
Synthetic human secretin did not improve symptoms or communication skills in children with autistic disorders .....	52		
		<b>Diagnosis</b>	
		The 15 item Geriatric Depression Scale had perfect sensitivity but low specificity for detecting depression in elderly people .....	55
		The Even Briefer Assessment Scale for Depression in German was accurate for identifying depression in elderly patients .....	56
		<b>Prognosis</b>	
		Dementia in advanced age led to higher mortality rates and shortened life .....	57
		<b>Aetiology</b>	
		Review: most antipsychotic drugs are associated with weight gain .....	58
		A lower minimum legal drinking age was associated with increased suicide in youths 18–23 years of age. ....	59
		The association between cognitive function and mortality pertained to specific but not general measures of cognitive function when health factors were considered .....	60
		Review: 7.7 to 12.9 units of alcohol per week is the level of alcohol consumption at which mortality is the lowest in men .....	61
		<b>Qualitative</b>	
		Resilience was not necessarily a healthy concept for adolescents in inner city vocational high schools. ....	62
		Physicians indicated the need to frame questions and develop indirect approaches that foster patient trust in evaluating victims of domestic violence .....	63
		St John's Wort users reported: a history of exposure to herbal remedies and belief in their safety; depressed mood; and ease of access to alternative medicines .....	64
		<b>Glossary</b> .....	Inside back cover