

<b>Supplementary Table 1. Proportion of missing data.</b>		
	Participants with SARS-CoV-2 infection prior to baseline (N = 770)	Participants without SARS-CoV-2 infection prior to baseline (N = 7654)
Gender	0.0%	0.0%
Age	0.0%	0.0%
Educational level	1.3%	1.0%
Household income	3.1%	4.5%
Tobacco smoking status	2.9%	3.2%
Body mass index	1.6%	1.4%
Self-rated health	0.7%	0.8%
Infection during follow-up	NA	0.0%
SSD-12 total score (at least one missing item)	22.6%	24.2%
SSD-12 item 1	0.6%	0.8%
SSD-12 item 2	0.8%	0.8%
SSD-12 item 3	0.7%	0.7%
SSD-12 item 4	0.8%	1.1%
SSD-12 item 5	0.7%	0.8%
SSD-12 item 6	0.8%	1.1%
SSD-12 item 7	19.8%	20.9%
SSD-12 item 8	1.6%	2.5%
SSD-12 item 9	1.1%	1.3%
SSD-12 item 10	4.6%	4.9%
SSD-12 item 11	1.8%	2.4%
SSD-12 item 12	1.1%	1.6%
Legends: Abbreviations: NA: not applicable; SSD-12: 12-item Somatic Symptoms Disorder-B Criteria Scale		